

NASPE Sets the Standard

POSITION STATEMENT

Comprehensive School Physical Activity Programs

The National Association for Sport and Physical Education (NASPE) recommends that all P-12 schools implement a Comprehensive School Physical Activity Program.

Schools play an important role in public health, and the physical, mental, and social benefits of regular physical activity for youth are well documented. Leading public health, medical, and educational organizations, including NASPE, have made important physical activity recommendations for school-aged youth (Centers for Disease Control and Prevention (CDC), 1997; Kaplan et al., 2005; NASBE, 2000; Pate et al. 2006; U. S. Department of Health and Human Services, 2000). These recommendations are for children to accumulate at least 60 minutes of physical activity on all or most all days of the week. (CDC, 2001; CSMF/CSH, 2006; Kaplan et al., 2005; Pate et al., 2006; NASPE, 2004; Strong et al., 2005; U.S. Department of Health and Human Services, 2004). In addition to physical activity recommendations, most of these agencies and organizations also include calls for schools to assume strong leadership roles in the education and promotion of physical activity among children (CDC, 1997; Kaplan et al., 2005; Pate et al., 2006; USDHHS, 1996; WHO, 2001). Conclusions drawn from the results of the 2006 School Health Policies and Programs Study (SHPPS) conducted by the Centers for Disease Control and Prevention include the need to implement a comprehensive approach at the state, district, and school levels to enhance physical education and physical activity in schools (Lee et al. 2007).

In 2004, federal legislation (PL 108-265) was passed which required all districts with federally funded school meal programs to develop and implement wellness policies by the beginning of the 2006-07 school year. The *Child Nutrition and WIC Reauthorization Act of 2004* directs school districts to set goals for physical activity, nutrition education, campus food provision, and other school-based activities designed to encourage student wellness. Furthermore, districts are required to engage a wide range of individuals in policy development and to have a plan for measuring policy implementation. Implementation of these wellness plans should result in an increase in school-based opportunities for physical activity.

A Comprehensive School Physical Activity Program (CSPAP) encompasses physical activity programming before, during, and after the school day. NASPE recommends that a CSPAP include: quality physical education; school-based physical activity

opportunities; school employee wellness and involvement; and family and community involvement.

Quality Physical Education

Quality physical education serves as the foundation of the CSPAP. According to NASPE (2004), a quality physical education program includes:

- Daily physical education (at least 150 minutes per week for elementary, and 225 minutes per week for middle/high school)
- A curriculum that meets the National Standards for Physical Education
- Student assessment aligned with instruction
- A certified physical education teacher providing meaningful content through standards-based instruction
- A pupil-teacher ratio equivalent to that in the classroom context
- Adequate equipment to promote maximum practice time

Research shows that quality physical education programs can contribute to students' regular participation in physical activity (Fairclough & Stratton, 2005; Luepker et al.,1996; Morgan, Beighle, Pangrazi, in press; NASPE, 2003; Sallis, McKenzie, Alcaraz, Kolody, Faucette, & Hovell, 1997), and can increase student participation in moderate to vigorous physical activity (CDC, 2001). Quality physical education is at the core of a CSPAP because of its role in helping students gain the knowledge and skills to become proficient movers and participants in a lifetime of physical activity (Baranowski et al., 1997; NASPE, 2004). Within the CSPAP, certified physical educators not only teach physical education classes, but also serve as physical activity leaders in their schools and promote physical activity both within and beyond the regular school day.

School-based Physical Activity Opportunities

Along with physical education classes, students need physical activity opportunities throughout the school day to meet the recommended minimum requirements of at least 60 minutes of physical activity each day. These physical activity opportunities are not to take the place of physical education, but rather supplement physical activity time accumulated during physical education class, and use the skills and knowledge learned in physical education to successfully be physically active.

During the school day, children and youth need a "break" from sedentary activities in the classroom. Physical activity breaks meet this need and can increase individuals' daily physical activity levels. Integrating physical activity programs in the school day has not been shown to decrease academic achievement. In fact, in some cases it may even enhance academic performance (Dywer, 1983; Sallis, McKenzie, Kolody et al., 1999; Shephard, LaVallee, Volle, LaBarre, & Beaucage, 1994; Shephard, 1997). Sedentary

classroom activities should be interspersed with bouts of physical activity, and when possible have movement integrated into the academic content. Classroom physical activity helps to mediate often cited barriers to learning such as inattentiveness and misbehavior (Mahar et al., 2006; Pelligrini et al., 1995).

For elementary school students, recess can be utilized to provide opportunities for students to accumulate meaningful amounts of physical activity (Beighle, Morgan, Le Masurier, & Pangrazi, 2006; Jago & Baranowski, 2004; Johns & Ha, 1999; Mota et al., 2005; Ridgers & Stratton, 2005). NASPE's position statement on Recess for Elementary School Students states that all elementary school children should be provided with at least one daily period of recess for a minimum of 20 minutes, and provides information on the importance of recess for a child's physical, social, and academic development (NASPE, 2006). In middle and high schools, students might spend free time, such as during lunch or study hall, participating in "drop in" physical activity sessions. For example, students might visit a supervised fitness center or check out a pedometer to monitor steps walked on campus during an unencumbered period.

Physical activity opportunities and programs occurring before and after the regular school hours, including intramural and interscholastic programs, have great potential for increasing overall daily physical activity levels of youth (Allison & Adlaf, 2000; Kaplan et al.,2005; NASPE, 2002a; NASPE, 2002b; Powers et al.,2002; Yin et al.,2005; Young et al.,2007), and should be included in a CSPAP. Intramural programs are those offered within the school environment and should provide a diverse selection of physical activity opportunities that meet the needs, interests, and abilities of most students. All students should be given the opportunity to participate in intramural activities, and special consideration should be given to those with particular physical activity needs and who are at-risk for a sedentary lifestyle (e.g., children with physical and other disabilities, those with chronic illnesses, those who are overweight).

Intramural activities can include sports (e.g., volleyball and basketball), self-directed activities (e.g., walking and jogging), classes (e.g., dance, yoga, or martial arts), and activity clubs (e.g., jump rope, hiking, and fitness). These activities can be offered before school begins, as well as after school. Other before-school activities designed to increase physical activity might include large group energizing sessions and/or a variety of small group physical activities from which students can choose.

Interscholastic sports programs are an important part of the American culture and provide physical activity opportunities for the more skillful students, particularly in middle and high schools. They should be offered in addition to school intramural programs.

Students and staff are encouraged to take active transport to school, such as walking and biking. Students who walk or bike to school generally expend more energy overall throughout the day (Tudor-Locke, Neff, Ainsworth, Addy, & Popkins, 2002; Tudor-Locke, Ainsworth, Adair, & Popkin, 2003) and this may particularly help overweight students (Rosenberg et al., 2006).

In providing school-based physical activity opportunities, efforts should be made to stimulate interest and participation from the greatest proportion of students in the school. This requires offering a wide variety of activities including non-competitive and health-enhancing choices. Consideration should also be given to providing opportunities at various times so that participation is feasible for all children (Lounsbery, Bungum, & Smith, 2007).

School Employee Wellness and Involvement

School employee wellness programs as well as staff involvement in the CSPAP implementation are important to school-wide success. School employee wellness programs have been shown to improve staff health, increase physical activity levels, and be cost effective (Eaton, Marx, & Bowie, 2007). Therefore, schools should plan and implement activities, policies, and incentive programs for faculty and staff members to encourage participating in and modeling a healthy lifestyle that includes physical activity. When school leaders are personally committed to good health practices, they serve as positive role models and may show increased support for student participation in physical activity.

Strong administrative support and staff involvement in the CSPAP are critical to school program success. School staff can assume various roles, such as volunteer sponsor, planner, and supervisor of activity-based events and programs (e.g., playground, recess, after school teams and clubs). The support, interest, and availability of the staff is critical to effective implementation of a CSPAP. Incentives, including stipends for extra standard duties, to increase the number of teacher/coach volunteers should be considered to support efforts in the CSPAP. Orientation and training should be offered to all school staff to facilitate cooperation and collaboration in implementing all components of the CSPAP.

Family and Community Involvement

The role of the school in promoting physical activity should extend beyond the school campus to reach families and collaborate with community members. Parents are an integral part of students' lives and can impact their physical activity levels (Nader et al., 1996; Ornelas, Perreira, & Ayala, 2007; Welk, Wood & Morss, 2003). Therefore, activities and events that include families, before, during, or outside the school day, are encouraged. Examples include family fun/fitness nights, after school classes for families, and use of facilities for family active participation after school and on weekends. Parents should be provided with information on the importance of physical activity and with strategies to promote their children's physical activity opportunities outside of school. This might include sharing information about physical activity and physical education on a regular basis through a variety of written, web-based, and event media outlets.

Additional physical activity opportunities can be made available to students through school collaboration with community-based providers of physical activity (Jago & Baranowski, 2004; Stevens et al., 2005). These collaborations could include: the use of school facilities for community recreation; the use of community facilities and programs

for promoting student/family physical activity; and the sharing of personnel and other resources to help provide more PA opportunities for all (Pate et al., 2006). Collaborative efforts to provide transportation from school to community facilities can increase student participation.

Action Statement

To fully establish a Comprehensive School Physical Activity Program, NASPE recommends that schools implement the following steps:

- 1. Appoint a committee to develop and oversee the CSPAP. This committee, perhaps a sub-committee of the School Wellness Policy Committee, should be comprised of (but not limited to) physical educators, health teachers, other teachers, administrators, parents, students, community members, and other stakeholders who can help facilitate the school's plan.
- 2. Conduct a baseline assessment of the CSPAP components as they currently exist in the school.
- 3. Create a vision statement and action plan appropriate to the specific school, addressing each of the four components of a CSPAP.
 - a. The action plan should include the baseline assessment, objectives. activities to meet each objective, defined outcomes, timelines, and persons responsible for each CSPAP component area:
 - Quality physical education
 - Before-school strategies
 - During-school strategies (outside of physical education)
 - After-school strategies
 - Staff personal wellness
 - School employee wellness and involvement in the school's CSPAP
 - Family and community involvement
- 4. Implement the Action Plan

References

Allison, K.R., & Adlaf, A.M. (2000). Structured opportunities for student physical activity in Ontario elementary and secondary schools. Canadian Journal of Public Health. 91(5), 371-375.

Baranowski, T., Bar-Or, O., Blair, S., Corbin, C., Dowda, M., Freedson, R., Pate, R., Plowman, S., Sallis, J., Saunders, R., Seefeldt, V., Siedentop, D., Simons-Morton, B., Spain, C., Tappe, M., & Ward D. (1997). Guidelines for school and community programs to promote lifelong physical activity among young people. *Morbidity and* Mortality Weekly Report 50, RR-6: 1-36.

- Beighle, A., Morgan, C.F., Le Masurier, G., & Pangrazi, R. P. (2006). Children's physical activity during recess and outside of school. Journal of School Health, *76*(10), 516-520.
- Centers for Disease Control and Prevention. (1997). "Guidelines for School and Community Programs to Promote Lifelong Physical Activity among Young People," Morbidity and Mortality Weekly Report 46, RR-6: 1-36, available online at: www.cdc.gov/HealthyYouth/physicalactivity/guidelines.
- Centers for Disease Control and Prevention. (2001). "Increasing physical activity: A report on recommendations of the task force on community prevention services," Morbidity and Mortality Weekly Report 50, RR-18: 1-16.
- Council on Sports Medicine and Fitness and Council on School Health. (2006). Active healthy living: Prevention of childhood obesity through increased physical activity. Pediatrics, 117, 1834-1842.
- Dwyer, T., Coonan, W., Leitch, D., Hetzel, B., & Baghurst, R. (1983). An investigation of the effects of daily physical activity on the health of primary school students in South Australia. International Journal of Epidemiologists, 12(3), 308-313.
- Eaton, D.K., Marx, E., & Bowie, S.E. (2007). Faculty and staff health promotion: Results from the school health policies and programs study 2006. Journal of School Health, 77(8), 557-566.
- Fairclough, S., & Stratton, G. (2005), Physical education makes you fit and healthy: Physical education's contribution to young people's physical activity levels. *Health* Education Research, 20(1), 14-23.
- Jago, R., & Baranowski, T. (2004). Non-curricular approaches for increasing physical activity in youth: A review. Preventive Medicine, 39(1), 157-163.
- Johns, D.P., & Ha, A.S. (1999). Home and recess physical activity of Hong Kong children. Research Quarterly for Exercise and Sport, 70(3), 319-323.
- Kaplan, J.P., Liverman, C.T., & Kraak, V.A.; (Eds.) (2005). Preventing childhood obesity: Health in the balance (pp.237-284). Washington, DC: The National Academies Press.
- Lee, S., Burgeson, C., Fulton, J., & Spain, C. (2007). Physical Education and physical activity: Results from the School Health Policies and Programs Study 2006. Journal of School Health, 7(1), 435-463.
- Luepker, R.V., Perry, C.L., McKinlay, S.M., Nader, P.R., Parcel, G.S, Stone, E.J. et al. (1996). Outcomes of a field trial to improve children's dietary patterns and physical

Comprehensive School Physical Activity Programs (Cont.)

- activity: The child and adolescent trial for cardiovascular health. Journal of the American Medical Association, 275(10), 767-776.
- Lounsbery, M., Burgum, T., & Smith, N. (2007). Physical activity opportunity in Nevada K-12 public school settings. Journal of Physical Activity and Health, 4(1), 30-38.
- Mahar, M.T., Murphy, S.K., Rowe, D.A., Golden, J., Shields, A.T., & Raedeke, T.D. (2006). Effects of a classroom-based program on physical activity and on-task behavior. Medicine & Science in Sports and Exercise, 38(12),286-294.
- Model School Wellness Policies. National Alliance for Nutrition and Activity. Available at: http://www.schoolwellnesspolicies.org. Accessed June 25, 2007.
- Morgan, C.F., Beighle, A., & Pangrazi, R.P. (2007). What are the contributory and compensatory relationships between physical education and physical activity in children? Research Quarterly in Exercise and Sport, 78(5), 407-412.
- Mota, J., Silva, P., Santos, M.P., Ribeiro, J.C., Oliverira, J., & Duarte, J.A. (2005). Physical activity and school recess time: Differences between the sexes and the relationship between children's playground physical activity and habitual physical activity. Journal of Sport Science, 23(3), 269-275.
- Nader, P.R., Sellers, D.E., Johnson, C.C., Perry, C.L., Stone, E.J., Cook, K.C., et al. (1996). The effect of adult participation in a school-based family intervention to improve children's diet and physical activity: The child and adolescent trial for cardiovascular health. Preventive Medicine, 25, 455-464
- National Association for Sport and Physical Education. (2002a). Co-curricular physical activity and sport programs for middle school students. Reston, VA: Author.
- National Association for Sport and Physical Education. (2002b). Guidelines for after school physical activity and intramural sport programs. Reston, VA: Author.
- National Association for Sport and Physical Education. (2004). Physical activity for children: A statement of guidelines for children ages 5-12. Reston, VA: Author.
- National Association for Sport and Physical Education. (2006). Recess for elementary school students. Reston, VA: Author.
- National Association for Sport and Physical Education. (2003). What constitutes a quality physical education program? Reston, VA: Author.
- National Association of State Boards of Education. (2000). Fit, Healthy and Ready to Learn: A School Health Policy Guide. Part 1: Physical Activity, Healthy Eating, and Tobacco Use Prevention. (Alexandria, VA: Author)

- Ornelas, I.J., Perreira, K.M., & Ayala, G.X. (2007). Parental influences on adolescent physical activity: A longitudinal study. International Journal of Behavioral Nutrition and Physical Activity, 4:3.
- Pate, R.R., Davis, M.G., Robins, T.N., Stone, E.J., McKenzie, T.L., & Young, J.C. (2006). Promoting physical activity in children and youth: A leadership role for schools. Circulation: Journal of the American Heart Association. 114, 1214-1224.
- Pellegrini, A.D., Huberty, P.D., & Jones, I. (1995). The effects of recess timing on children's playground and classroom behaviors. American Educational Research Journal, 32(4), 845-864.
- Powers, H.S., Conway, T.L., McKenzie, T.L., Sallis, J.F., & Marshall, S.J. (2002). Participation in extracurricular physical activity programs at middle schools. Research Quarterly for Exercise and Sport, 73(2), 187-192.
- Ridgers, N.D., & Stratton, G. (2005). Physical activity during school recess: The Liverpool sporting playgrounds project. Pediatric Exercise Science, 17, 281-290.
- Rosenberg, D.E., Sallis, J.F., Conway, T.L., Cain, K.L., & McKenzie, T. L. (2006). Active transportation to school over 2 years in relation to weight status and physical activity. Obesity, 14(10), 1771-1776.
- Sallis, J.F., McKenzie, T.L., Alcaraz, J.E., Kolody, B., Faucette, N., & Hovell, M.F. (1997). The effects of a 2-year physical education program (SPARK) on physical activity and fitness in elementary school students. American Journal of Public Health, 87, 1328-1334.
- Sallis, J.F., McKenzie, T.L., Kolody, B., Lewis, M., Marshall, S., & Rosengard, P. (1999). Effects of health-related physical education on academic achievement: Project SPARK. Research Quarterly in Sport and Exercise, 70(2), 127-134.
- Shephard, R.J., LaVallee, H., Volle, M., LaBarre, R., & Beaucage, C. (1994). Academic skills and required physical education: The Trois Rivieres experience. CAHPER Research Supplement, 1(1), 1-12.
- Shephard, R.J. (1997). Curricular physical activity and academic performance. *Pediatric* Exercise Science, 9, 113-126.
- Stevens, J., Murray, D.M., Catellier, D.J., Hannan, P.J., Lytle, L.A., Elder, J.P., Yong, et al. (2005). Design of the trial of activity for adolescent girls (TAGG). Contemporary Clinical Trials, 26: 223-233.
- Strong, W.B., Malina, R.M., Blimkie, C.J.R., Daniels, S.R., Dishman, R.K., Gutin, B., et al.(2005). Evidence based physical activity for school-age youth. The Journal of Pediatrics, 146(6), 732-737.

- Timperio, A., Ball, K., Salmon, J., Roberts, R., Giles-Corti, B., Simmons, D., Baur, L.A., & Crawford, D. (2006). Personal, family, social, and environmental correlates of active commuting to school. American Journal of Preventive Medicine, 30(1), 45-51.
- Tudor-Locke, C., Ainsworth, B.E., Adair, L.S., & Popkin, B.M. (2003). Objective physical activity of Filipino youth stratified for commuting mode to school. Medicine & Science in Sports and Exercise, 35, 465-471.
- Tudor-Locke, C., Neff., L.J., Ainsworth, B.E., Addy, C.L., & Popkin, B.M. (2002). Omission of active commuting to school and the prevalence of children's healthrelated physical activity levels: The Russian longitudinal monitoring study. Child: Care, Health & Development, 28, 507-512.
- U.S. Department of Health and Human Services. (2000). Healthy People 2010 (Conference Edition, in Two Volumes). Washington, DC: U.S. Government Printing Office.
- U.S. Department of Health and Human Services, U.S. Department of Agriculture. (2004). Dietary Guidelines for Americans, 2005. Washington, DC: U.S. Department of Health Services/U.S. Department of Agriculture.
- U.S. Department of Health and Human Services (USDHHS). (1996). *Physical Activity* and Health: A Report of the Surgeon General. Atlanta: Centers for Disease Control and Prevention.
- Welk, G.J., Wood, K., & Morss, G. (2003). Parental influences on physical activity in children: An exploration of potential mechanisms. Pediatric Exercise Science, 15, 19-33.
- World Health Organization. (2001). Obesity: Preventing and managing the global epidemic. Report of a WHO Consultation on Obesity, 3-5 June 1997, Geneva. Geneva, Switzerland: World Health Organization. WHO/NUT/NCD 98.1
- Yin, Z., Moore, J.B., Johnson, M.H., Barbeau, P., Cavnar, M., Thornburg, J., & Gutin, B. (2005). The Medical College of Georgia FitKid Project: The relations between program attendance and changes in outcomes in year 1. International Journal of Obesity, 29, S40-S45.
- Young, D.R., Felton, G.M., Grieser, M., Elder, J.P., Johnson, C., Lee, J., et al. (2007). Policies and opportunities for physical activity in middle school environments. Journal of School Health, 77(1), 41-47.

Comprehensive School Physical Activity Programs Task Force

Aaron Beighle, University of Kentucky Eloise Elliott, Concord University, WV Thom McKenzie, San Diego State University, CA Amelia Mays Woods, University of Illinois at Urbana-Champaign

Funding for this position statement was made possible in part by cooperative agreement award number U58/DP-000420 from the Centers for Disease Control and Prevention.

The views expressed in written presentation materials and by speakers do not necessarily reflect the official policies of the Department of Health and Human Services; nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.

Suggested Citation:

National Association for Sport and Physical Education. (2008). Comprehensive school physical activity programs [Position statement]. Reston, VA: Author.